

MEASURING σ_8 WITH CLUSTER LENSING : BIASES FROM UNRELAXED CLUSTERS

GRAHAM P. SMITH,^{1,2} ALASTAIR C. EDGE,³ VINCENT R. EKE,³ ROBERT C. NICHOL,⁴ IAN SMAIL,³ &
 JEAN-PAUL KNEIB,^{2,5}

Draft version February 2, 2008

ABSTRACT

We use gravitational lens models and X-ray spectral analysis of ten X-ray luminous galaxy clusters at $z \simeq 0.2$ to study the impact of cluster substructure on attempts to normalize the matter power spectrum. We estimate that unrelaxed clusters are 30% hotter than relaxed clusters causing σ_8 to be over-estimated by 20% if the cluster selection function is not accounted for correctly. This helps to explain the wide range in σ_8 derived from different techniques, $\sigma_8 \sim 0.6$ – 1 , and offers a physically motivated explanation for some of the discrepancy. We also identify two further systematics in our analysis: (i) extrapolation of small field-of-view mass measurements to the cluster virial radius and (ii) projection of 3-dimensional masses contained in numerical simulations to the 2-dimensional information that is available from observations. We combine quantitative estimates of these two effects with our model fitting to estimate from the current data that $\sigma_8 = 0.75 \pm 0.05$ (statistical) ± 0.15 (systematic), where the systematic error reflects the extrapolation and projection uncertainties. All three systematics (substructure, extrapolation and projection) are fundamental to future cluster-based measurements of σ_8 regardless of the techniques employed. However, we identify gravitational lensing as the tool of choice for such studies, because a combination of strong- and weak-lensing offers the most direct route to control the systematics and thus achieve an unbiased comparison between observation and theory.

Subject headings: cosmology:observations — gravitational lensing — X-rays:galaxies:clusters — large-scale structure of the universe

1. INTRODUCTION

The spectrum of cosmic matter fluctuations is an important constraint on theoretical models of structure formation (e.g. Press & Schechter 1974; Bond et al. 1991; Bower 1991; Kauffmann & White 1993; Lacey & Cole 1993). The amplitude of the power spectrum is parametrized as σ_8 , the linear-theory value of the rms fractional fluctuations in density averaged in spheres of $8 h^{-1}$ Mpc radius at $z = 0$. Several methods have been used to estimate σ_8 : measurement of the abundance of galaxy clusters (Eke, Cole & Frenk 1996; Borgani et al. 2001; Reiprich & Böhringer 2001; Seljak 2001; Allen et al. 2002; Viana, Nichol & Liddle 2002; Pierpaoli et al. 2001), cosmic shear analyses (e.g. Bacon et al. 2002; Hoekstra et al. 2002; Refregier et al. 2002; van Waerbeke et al. 2002; Brown et al. 2002; Jarvis et al. 2002), cosmic microwave background (CMB) studies (Sievers et al. 2002; Bond et al. 2002), combined analysis of galaxy redshift survey and CMB data (Lahav et al. 2002). Current estimates of σ_8 range from ~ 0.6 to ~ 1.0 , with claimed statistical uncertainties in the range $\Delta \sigma_8 \sim 0.02$ – 0.15 . Overall, the situation is characterized by a lack of agreement between the results from different methods, or the same methods used on different samples, suggesting that systematic uncertainties probably lie at the heart of the current disagreement over the value of σ_8 .

In this letter we investigate systematic biases in the use of cluster abundances to measure σ_8 . In principal the mass function of galaxy clusters, $n(>M)$, should yield a direct constraint on σ_8 . However, it is not currently possible to measure cluster masses with the precision and in the numbers required to

construct a robust cluster mass function from direct measurement. The local cluster X-ray temperature function, $n(>T)$, has proved more accessible (e.g. Edge et al. 1990; Henry & Arnaud 1991; Markevitch 1998; Blanchard et al. 2000; Pierpaoli et al. 2000; Ikebe et al. 2002). The X-ray temperature function in conjunction with a robust mass-temperature calibration therefore offers an opportunity to constrain σ_8 .

Observational attempts to calibrate the mass-temperature relation typically rely on X-ray observations of clusters (e.g. Markevitch 1998; Nevalainen, Markevitch & Forman 2000; Finoguenov, Reiprich & Böhringer 2001; Allen, Schmidt & Fabian 2001, hereafter ASF; Reiprich & Böhringer 2002). Despite the progress made by Allen (1998) in understanding X-ray based cluster mass measurements, X-ray techniques continue to assume that all clusters are symmetric, equilibrium systems. This is a major concern, because ~ 40 – 70% of galaxy clusters appear to be dynamically immature (e.g. Mohr et al. 1995; Buote & Tsai 1996; Ota & Mitsuda 2002; Smith et al. 2003, in prep., hereafter S03), and this immaturity has a measurable systematic impact on the normalization of the cluster mass-temperature and mass-luminosity relations (Ota & Mitsuda 2002; S03; see also Randall et al. 2002).

In contrast, mass estimates based on gravitational lensing are insensitive to the physical nature and state of the cluster mass. Cluster lensing studies are therefore free from the symmetry and equilibrium assumptions that plague the X-ray studies. Attempts to use lensing to calibrate the cluster mass-temperature relation have so far relied on previously published and/or crude cluster mass estimates (Hjorth et al. 1998; ASF; Viana, Nichol

¹Department of Physics, University of Durham, South Road, Durham DH1 3LE, UK

²Current address: California Institute of Technology, MC 105-24, Pasadena, CA 91125, USA – Email: gps@astro.caltech.edu

³Institute for Computational Cosmology, Department of Physics, University of Durham, South Road, Durham DH1 3LE, UK

⁴Department of Physics, Carnegie Mellon University, 5000 Forbes Avenue, Pittsburgh, PA 15213-3890, USA

⁵Observatoire Midi-Pyrénées, 14 Avenue E. Belin, 31400 Toulouse, France

& Liddle 2002). A major improvement on these pioneering studies would come from a precise and uniform analysis of a large objectively selected cluster sample for which high resolution space-based optical and X-ray data were available. In anticipation of such a program we conduct a pilot study using S03's *Hubble Space Telescope (HST)/Chandra* gravitational lensing survey of ten X-ray luminous galaxy clusters at $z \simeq 0.2$. S03 made precise cluster mass and temperature measurements and thus constrained the high-mass end of the cluster mass-temperature relation. They also studied the dependence of this normalization on cluster substructure, concluding that unrelaxed clusters are, on average, 30% hotter than relaxed clusters. S03's results therefore offer a unique opportunity to study the impact of cluster substructure on estimates of σ_8 .

We summarize S03's results in §2, describe our modeling and results in §3 and summarize our conclusions in §4. We express the Hubble parameter in terms of h , where $H_0 = 100 h \text{ km s}^{-1} \text{ Mpc}^{-1}$. We also adopt $\Omega_0 = 0.3$ and $\Lambda_0 = 0.7$.

2. HST/CHANDRA MASS-TEMPERATURE CALIBRATION

S03 studied a representative sample of ten of the most X-ray luminous clusters ($L_X \geq 2 \times 10^{44} h^{-2} \text{ erg s}^{-1}$, 0.1–2.4 keV) in a narrow redshift slice at $z = 0.21 \pm 0.04$, with line-of-sight reddening of $E(B-V) \leq 0.1$ from the XBACs sample (Ebeling et al. 1996). Each cluster was typically observed for 3 orbits (i.e. 7.5ks) through the F702W filter using the WFPC2 camera on-board *HST*. S03 used these data in conjunction with ground-based optical and near-infrared data (Smith et al. 2001, 2002), and the LENSTOOL software (Kneib 1993; Kneib et al. 1996; Smith 2002) to construct a detailed gravitational lens model of each cluster.

Armed with these models, S03 measured M_{2500} , the total projected cluster mass within r_{2500} , i.e. the radius at which the density of matter in the clusters falls to $\rho = \rho_{2500} = 2500 \rho_c$, where ρ_c is the critical density required to close the universe.⁶ S03 also used the models to divide their sample into relaxed ($M_{\text{sub}}/M_{\text{tot}} < 10\%$) and unrelaxed ($M_{\text{sub}}/M_{\text{tot}} > 10\%$) clusters where M_{tot} is the total projected mass of the cluster within r_{2500} and M_{sub} is the projected mass of the cluster within the same radius that is not associated with the main centrally-located dark matter halo and cluster central galaxy. A complementary analysis of archival *Chandra* and *ASCA* observations of eight and one of these clusters respectively also provided accurate measurements of the temperature of each cluster ($T_{X,\text{tot}}$) within a projected radius of $r \leq 1 h^{-1} \text{ Mpc}$. We refer the reader to S03 for further details of the modeling and analysis of these clusters.

We plot S03's mass and temperature measurements in Fig. 1. The open symbols show the individual clusters, and the filled symbols indicate the properties of the mean relaxed and unrelaxed cluster sub-samples. The mean temperatures of the relaxed and unrelaxed clusters, are $\langle T_{X,\text{tot}} \rangle = 6.3 \pm 0.8 \text{ keV}$ and $\langle T_{X,\text{tot}} \rangle = 9.2 \pm 1.2 \text{ keV}$ respectively, where the error bars are the uncertainties on the means and are the estimated from bootstrap re-sampling each sub-sample of clusters; we therefore expect these error bars to be over-estimates. The unrelaxed clusters appear to be systematically hotter than the relaxed clusters.

Two of S03's sample (A383 – Smith et al. 2001; A1835 – e.g. Schmidt et al. 2001) have central cooling timescales of $t_{\text{cool}} \lesssim 10^9 \text{ years}$. This is in line with expectations from other representative samples of X-ray luminous clusters (Peres et al. 1998). S03 therefore recalculated all of the cluster tempera-

tures using an $0.05 \leq r \leq 1 h^{-1} \text{ Mpc}$ annulus (i.e. excluding the cold core of the two extreme “cooling flow” systems). They found that, while the temperature difference is slightly reduced ($\langle T_{X,\text{ann}} \rangle_{\text{relaxed}} = 6.9 \pm 0.9 \text{ keV}$), it is robust to the exclusion of the central $50 h^{-1} \text{ kpc}$ of each cluster from the temperature calculations. The 30% temperature difference therefore does reflect a bona fide difference between the ambient temperatures of relaxed and unrelaxed clusters. We note that this affect is similar to a substructure-related bias found by Ota & Mitsuda (2002) in the cluster mass-luminosity relation.




FIG. 1. — M_{2500} , the projected mass within r_{2500} , versus the temperature of the intra-cluster medium for S03's sample of X-ray luminous clusters. The unrelaxed clusters are on average 30% hotter than the relaxed clusters, causing a previously unquantified structural bias in the normalization of the cluster mass-temperature relation. We also plot S03's and ASF's mass-temperature relations, assuming a canonical logarithmic slope of $\alpha = 2/3$ for the S03 relation. The ASF relation agrees with the two cooling flow clusters in S03's sample (A383 and A1835 are the two open circles that lie within $1-\sigma$ of the line).

3. MODELING AND RESULTS

3.1. Approach

We construct a simple model to investigate the impact of S03's results on estimates of σ_8 . We begin by parameterizing the cluster mass-temperature relation:

$$T_{X,\text{tot}} (\text{keV}) = A(M_{2500}/10^{14} h^{-1} M_{\odot})^{\alpha} \quad (1)$$

where $T_{X,\text{tot}}$ and M_{2500} are as defined in §2, and A, α are the normalization and logarithmic slope respectively. We first convert the Jenkins et al. (2001) mass function to a temperature function. This conversion includes the following elements: a mass-dependent concentration index (Eke, Navarro & Steinmetz 2001), conversion of three-dimensional masses from the simulations to projected two-dimensional masses (Hjorth et al. 1998) and an observational mass-temperature normalization (S03, ASF). We then fit this model temperature function to the observed temperature function (Edge et al. 1990) using a single free parameter, σ_8 .

Our model also contains the following parameters: $\{\Omega_0, \Lambda_0, \Gamma, \sigma_T\}$ where Ω_0 and Λ_0 are the matter and vacuum energy

⁶At $z = 0.2$, r_{2500} corresponds to the edge of the *HST*/WFPC2 field of view for the most massive clusters in S03's sample.

densities of the Universe at $z = 0$, Γ is the spectral shape parameter for the power spectrum and σ_T is the scatter in $\log(T_{X,\text{tot}})$. We focus our attention on the dependence of σ_8 on A and to a lesser extent on σ_T ; we therefore adopt “concordance” values for the remaining parameters: $\Omega_0 = 0.3$, $\Lambda_0 = 0.7$, $\Gamma = 0.2$ (e.g. Efstathiou et al. 2002), $\alpha = 2/3$ (e.g. ASF). We stress that we adopt a fixed value of $\Omega_0 = 0.3$, and therefore do not investigate the σ_8 – Ω_0 degeneracy.

3.2. Model Fitting

We use two independent mass-temperature calibrations to normalize our models. We begin with S03’s normalization, and adopt the values of A and σ_T relevant to their entire sample: $A = 4.4$, $\sigma_T = 0.1$ (see solid line in Fig. 1). We compute a model temperature function and fit it to the observed temperature function (Edge et al. 1990), obtaining a best-fit of $\sigma_8 = 0.75 \pm 0.05$ where the uncertainty is the statistical uncertainty on the fit. We plot this best-fit model and the observed temperature function in Fig. 2. Next, we turn to ASF’s cooling flow mass-temperature relation. These authors observed a sample of seven cooling flow clusters with *Chandra*, and used these data to normalize the mass-temperature relation. We convert ASF’s cooling flow mass-temperature relation into the form required for our model: $A = 2.6$, $\sigma_T \simeq 0.03$. Using these values, we construct a model temperature function and fit it to the observed function, obtaining $\sigma_8 = 0.91 \pm 0.07$. This model (Fig. 2) fits the data less well than the S03-based model, with the largest residuals occurring at high temperatures.




FIG. 2. — We plot the Edge et al. (1990) cluster temperature function for both all clusters and cooling flow clusters (defined as containing a line-emitting central galaxy), together with the best-fit model temperature functions that are normalized with the S03 and ASF mass-temperature relations. When a cooling flow cluster based normalization is applied to a representative sample of clusters, σ_8 is over-estimated by $\sim 20\%$. However, when the cluster selection function is accounted for properly in both the model normalization and the observed temperature function, consistent values of σ_8 are obtained.

A simple interpretation of these two models is that it is important to understand the cluster selection function when using cluster abundances to measure σ_8 . Specifically, using a cooling-flow cluster mass-temperature normalization when studying the

temperature function of a representative sample of clusters may cause σ_8 to be over-estimated by approximately 20%.

We test this interpretation by fitting the cooling flow normalized model to a temperature function that describes just cooling flow clusters. We first use the observed correlation between line emission from cluster central galaxies and short cooling timescales ($t_{\text{cool}} \lesssim 10^9$ years – Edge et al. 1992; Peres et al. 1998; Crawford et al. 1999) to construct a “cooling flow only” temperature function from the Edge et al. (1990) sample. We then fit the cooling flow model to the cooling flow data and obtain a best-fit value of $\sigma_8 = 0.74 \pm 0.05$, which agrees with the value obtained from the original model that was normalized with S03’s results. We plot this best-fit model and the relevant data in Fig. 2. This model confirms our interpretation that cluster substructure is an important and previously unidentified systematic effect at the 20% level when using cluster abundances to constrain σ_8 .

3.3. Further Uncertainties

The low value of σ_8 obtained in §3.2 is similar to a number of other recent results that favor $\sigma_8 \sim 0.6$ – 0.8 (e.g. Seljak 2001; Reiprich & Böhringer 2001; Borgani et al. 2001; Allen et al. 2002; Viana, Nichol & Liddle 2002; Lahav et al. 2002; Brown et al. 2002; Jarvis et al. 2002). However, several other uncertainties need to be investigated before a reliable conclusion on the value of σ_8 from cluster abundance determinations can be drawn.

Firstly, we highlight the extrapolation of S03’s lens models from r_{2500} (i.e. approximately the edge of the *HST*/WFPC2 field of view at $z \sim 0.2$) to the cluster virial radii as a key systematic uncertainty in our analysis. Bardeau et al. (in prep.) investigate this effect in detail through their weak-shear analysis of panoramic ($28' \times 42'$) CFH12k *BRI*-band imaging of the S03 cluster sample. Prior to the completion of this wide-field analysis, we note that weak lensing analyses of individual clusters (e.g. King, Clowe & Schneider 2002) are unable to discriminate between isothermal ($\rho \propto r^{-2}$) and Navarro, Frenk & White (1997) ($\rho \propto r^{-3}$) profiles on large scales. We therefore exploit this lack of discriminatory power to make a conservative estimate of this systematic uncertainty. We integrate both profiles over the radial range $0.25 \lesssim r \lesssim 1.5 h^{-1} \text{Mpc}$ (i.e. the dynamic range over which we are extrapolating), and estimate that the uncertainty in profile shape introduces an uncertainty in the virial mass estimate for an individual cluster of $\sim 30\%$, which translates into an uncertainty in cluster temperature (assuming $M \propto T_X^{3/2}$) of $\sim 20\%$. This equates to a $\sim 10\%$ “extrapolation” systematic uncertainty in σ_8 .

Secondly, we identify the projection of three-dimensional cluster masses from numerical simulations to observed two-dimensional masses (§3.1) as a further source of systematic uncertainty. As Hjorth et al. (1998) discuss, the magnitude of this uncertainty depends on the slope of the cluster density profile at small radii. Recent observational results (Smith et al. 2001; Sand, Treu & Ellis 2002; Dahle, Hannestad & Sommer-Larsen 2002) indicate that there may be substantial intrinsic scatter in this slope, appearing to contradict theoretical claims for a universal profile (e.g. Navarro, Frenk & White 1997). Given these complications, we conservatively adopt a further 10% “projection” systematic uncertainty in σ_8 .

In summary, although S03’s detailed lens models allow the “substructure” systematic to be accounted for properly and (to first order) eliminated from our analysis, “extrapolation” and “projection” uncertainties combine to produce a $\sim 20\%$ sys-

tematic that we are unable to control with the current dataset.

4. SUMMARY AND DISCUSSION

We have used S03's substructure-dependent cluster mass-temperature normalization to investigate the impact of cluster substructure on estimates of σ_8 . We find that when a cooling flow cluster mass-temperature normalization is applied to the general cluster population, σ_8 is over-estimated by 20%. A clear understanding of the cluster selection function is therefore fundamental to attempts to constrain σ_8 with cluster abundances. The simple X-ray luminosity-limited selection of S03's sample (§2) enable us to account for this "substructure" systematic from our analysis and thus to estimate that $\sigma_8 = 0.75 \pm 0.05$ (statistical). However, before we conclude that $\sigma_8 = 0.75$, we highlight two further systematic effects which may bias our analysis: extrapolation of S03's small field-of-view lens models out to the cluster virial radii, and uncertainties in the relationship between three-dimensional mass information contained in numerical simulations and the two-dimensional mass information that is available from observations. We estimate conservatively that these effects combine to produce a further 20% systematic uncertainty, and therefore we conclude from the present data that $\sigma_8 = 0.75 \pm 0.05$ (statistical) ± 0.15 (systematic). We also note that the recently reported discrepancies between *XMM*- and *Chandra*-based cluster temperature measurements (Schmidt et al. 2001; Majerowicz et al. 2002; Markevitch 2002) may introduce further uncertainty into cluster abundance determinations of σ_8 .

Our 20% "substructure" systematic is similar to the discrepancy between the canonical value of $\sigma_8 \sim 0.9$ –1 (e.g. Eke, Cole & Frenk 1996; Pierpaoli et al. 2001; Bacon et al. 2002; Bond et al. 2002; Hoekstra et al. 2002; Refregier et al. 2002; van Waerbeke et al. 2002) and recent claims for $\sigma_8 \sim 0.6$ –0.8 (Seljak 2001; Reiprich & Böhringer 2001; Borgani et al. 2001; Allen et al. 2002; Brown et al. 2002; Jarvis et al. 2002; La-

hav et al. 2002; Schuecker et al. 2002; Viana, Nichol & Liddle 2002). Our results therefore offer a physically motivated explanation for some of this discrepancy. Independent confirmation of this comes from Randall et al.'s (2002) semi-analytic study of the effect of cluster mergers on the observed luminosity and temperature functions, and thus on the inferred cluster mass function. Randall et al. predict that cluster mergers boost the observed temperature function and can cause σ_8 to be over-estimated by 20% if hydrostatic equilibrium is assumed for non-equilibrium clusters, in agreement with our observational results.

All three systematics discussed in this letter (substructure, extrapolation and projection) affect the ability of cluster abundance techniques to measure σ_8 accurately, regardless of whether gravitational lensing or X-ray techniques are used to measure the cluster masses. However, the insensitivity of gravitational lensing to the physical nature and state of the cluster matter means that a combined strong- and weak-lensing study of a large, objectively selected sample of clusters should be the tool of choice for future cluster abundance studies.

ACKNOWLEDGMENTS

We thank Steve Allen, Andrew Benson, Harald Ebeling, Richard Ellis, Andy Fabian, Carlos Frenk, Andrew Liddle, Pasquale Mazzotta, Tommaso Treu and Pedro Viana for useful discussions and assistance. GPS acknowledges a postgraduate studentship from PPARC. ACE, VRE and IRS acknowledge University Research Fellowships from the Royal Society. RCN thanks the Department of Physics at the University of Durham for their hospitality during the summer of 2002 when this work was performed, and is grateful for financial support under grant number NAG-5606. IRS acknowledges a Philip Leverhulme Prize Fellowship. JPK acknowledges financial support from CNRS.

REFERENCES

- Allen S.W., 1998, MNRAS, 296, 392
 Allen S.W., Schmidt R.W., Fabian A.C., 2001, MNRAS, 328, L37
 Allen S.W., et al., 2002, astro-ph/0208394
 Bacon D.J., et al., 2002, MNRAS, submitted, astro-ph/0203134
 Bond J.R., et al., 1991, ApJ, 379, 440
 Bond J.R., et al., 2002, ApJ, submitted, astro-ph/0205386
 Borgani S., Rosati P., Tozzi P., Stanford S.A., Eisenhardt P.R., Lidman C., Holden B., Della Ceca R., Norman C., Squires G., 2001, ApJ, 561, 13
 Bower R.G., 1991, MNRAS, 248, 332
 Brown M.L., et al., 2002, MNRAS submitted, astro-ph/0210213
 Buote D.A. & Tsai J.C., 1996, ApJ, 458, 27
 Crawford, C.S., et al., 1999, MNRAS, 306, 857
 Dahle H., Hannestad S., Sommer-Larsen J., 2002, ApJL, submitted, astro-ph/0206455
 Ebeling H., et al., 1996, MNRAS, 281, 799
 Edge A.C., Stewart G.C., Fabian A.C., Arnaud K.A., 1990, MNRAS, 245, 559
 Edge A.C., Stewart G.C., Fabian A.C., 1992, MNRAS, 258, 177
 Efstathiou G., Moody S., Glazebrook K. & the 2FGRS Team, 1996, Cole S., Frenk C.S., 1996, MNRAS, 282, 263
 Eke V.R., Navarro J.F., Steinmetz, 2001, 554, 114
 Finoguenov A., Reiprich T.H., Böhringer H., 2001, A&A, 368, 749
 Henry J.P. & Arnaud K.A., 1991, ApJ, 372, 410
 Hjorth J., Oukbir J., van Kampen E., 1998, MNRAS, 298, L1
 Hoekstra H., Yee H.K.C., Gladders M.D., et al., 2002, ApJ, 572, 55
 Ikebe Y., Reiprich T.H., Böhringer H., Tanaka Y., Kitayama T., 2002, A&A, 383, 773
 Jenkins A., Frenk C.S., White C.S., White S.D.M., Colberg J.M., Cole S., Evrard A.E., Couchman H.M.P., Yoshida N., 2001, MNRAS, 321, 372
 Kauffmann G. & White S.D.M., 1993, 261, 921
 King L.J., Clowe D.I., Schneider P., 2002, A&A, 383, 118
 Kneib J.-P., 1993, PhD Thesis, Université Paul Sabatier, Toulouse, France
 Kneib J.-P., Ellis R.S., Smail I., Couch W.J., Sharples R.M., 1996, ApJ, 471, 643
 Lacey C.G., Cole S., 1993, MNRAS, 262, 627
 Lahav O., Bridle S.L. & the 2dFGRS Team, 2002, MNRAS, 333, 961
 Majerowicz S., Neumann D.M., Reiprich T.H., 2002, A&A, submitted, astro-ph/0202347
 Markevitch M., 1998, ApJ, 504, 27
 Markevitch M., et al., 2000, ApJ, 541, 542
 Markevitch M., 2002, astro-ph/0205333
 Mohr J.J., Evrard A.E., Fabricant D.G., Geller M.J., 1995, ApJ, 447, 8
 Navarro, J.F., Frenk, C.S., White, S.D.M., 1997, ApJ, 490, 493
 Nevalainen J., Markevitch M., Forman W., 2000, ApJ, 532, 694
 Ota N. & Mitsuda K., 2002, ApJ, 567, L23
 Peres, C.B., Fabian, A.C., Edge, A.C., Allen, S.W., Johnstone, R.M., White, D.A., 1998, MNRAS, 298, 416
 Pierpaoli E., Scott D., White M., 2001, MNRAS, 325, 77
 Press W.H., Schechter P., 1974, ApJ, 187, 425
 Randall S.W., Sarazin C.L., Ricker P.M., 2002, ApJ, 577, in press (astro-ph/0206161)
 Refregier A., Rhodes J., Groth E.J., 2002, ApJ, 572, L131
 Reiprich T.H., Böhringer H., 2002, ApJ, 567, 716
 Sand D.J., Treu T., Ellis R.S., 2002, ApJ, 574, L129
 Schuecker P., Böhringer H., Collins C.A., Guzzo L., 2002, A&A, submitted, astro-ph/0208251
 Schmidt R.W., Allen S.W., Fabian A.C., 2001, MNRAS, 327, 1057
 Seljak U., 2001, MNRAS, submitted, astro-ph/0111362
 Sievers J.L., Bond J.R., Cartwright J.K., et al., ApJ, submitted, astro-ph/0205387
 Smith G.P., 2002, PhD Thesis, University of Durham, UK
 Smith G.P., Kneib J.-P., Ebeling H., Czoske O., Smail I., 2001, ApJ, 552, 493
 Smith G.P., Smail I., Kneib J.-P., Czoske O., Ebeling H., Edge A.C., Pello R., Ivison R.J., Packham C., Le Borgne J.-F., 2002, MNRAS, 330, 1
 van Waerbeke L., Mellier Y., Pello R., et al., 2002, A&A, submitted, astro-ph/0202503
 Viana P.T.P., Nichol R.C., Liddle A.R., 2002, ApJ, 569, L75